

Pulsar rotation measures and the magnetic structure of our Galaxy

J. L. Han^{1,2}, R. N. Manchester³, and G. J. Qiao^{2,4,5}

¹*Beijing Astronomical Observatory, Chinese Academy of Sciences (CAS), Beijing 100080, China*

²*Beijing Astrophysical Center, CAS-PKU, Beijing 100871, China*

³*Australia Telescope National Facility, CSIRO, PO Box 76, Epping, NSW 2121, Australia*

⁴*CCAT (World Laboratory), PO Box 8730, Beijing 100080, China*

⁵*Department of Geophysics, Peking University (PKU), Beijing 100871, China*

Accepted Feb.3, 1999; Received Jan.20, 1999; in original form 1997 November 6

ABSTRACT

We have obtained 63 rotation measures (RMs) from polarization observations of southern pulsars, of which 54 are new measurements and 3 are varied from previous values. The new pulsar RM data at high Galactic latitudes are mostly consistent with the antisymmetric RM distribution found previously. For the Galactic disc, evidence for a field reversal near the Perseus arm, and possibly another beyond it, is presented. Inside the Solar Circle, in addition to the two known field reversals in or near the Carina-Sagittarius arm and the Crux-Scutum arm, a further reversal in the Norma arm is tentatively identified. These reversals, together with the pitch angle derived from pulsar RM and stellar polarization distributions, are consistent with bisymmetric spiral (BSS) models for the large-scale magnetic field structure in the disc of our Galaxy. However, discrimination between models is complicated by the presence of smaller-scale irregularities in the magnetic field, as well as uncertainties in the theoretical modelling.

Key words: ISM: magnetic fields — Galaxy: structure — Galaxies: magnetic field — Pulsars: general

1 INTRODUCTION

Early observations of starlight polarization (eg. Mathewson & Ford 1970), the rotation measure (RM) distribution of extragalactic radio sources (e.g. Gardner, Morris & Whiteoak 1969; Simard-Normandin & Kronberg 1980) and pulsar RMs (Manchester 1974) revealed many of the principal features of the Galactic magnetic field. See Han & Qiao (1994) for a comprehensive review of earlier work.

Pulsars have a number of advantages as probes of the interstellar magnetic field. They typically have substantial linear polarization, making the Faraday rotation relatively easy to measure; they are distributed in the Galaxy with approximately known distances, allowing the three-dimensional properties of the magnetic field to be investigated; and they apparently have no intrinsic rotation measure (Manchester 1971). With the dispersion measure (DM), direct estimates of the Galactic magnetic field strength, averaged over the line of sight to the pulsar with a weight proportional to the local free electron density, can be obtained from

$$\langle B_{\parallel} \rangle = 1.232 \text{ RM/DM} \quad (1)$$

where $B_{\parallel}$ is in μG and RM and DM are in their usual units, rad m^{-2} and $\text{cm}^{-3} \text{ pc}$, respectively. For a positive RM, the

mean field is directed toward us. Manchester (1974) used RMs of 38 pulsars to estimate the strength and direction of the local Galactic magnetic field, concluding that the local field is basically azimuthal. Thomson & Nelson (1980) analysed the RMs of 48 pulsars, finding evidence for a field reversal in the inner Carina-Sagittarius spiral arm.

More extensive studies became possible only after the RMs of a large sample of pulsars had been measured. Lyne & Smith (1989) analysed the RMs of 185 pulsars (Hamilton & Lyne 1987), and found evidence for a reversal about 1 kpc inside the Solar Circle. Comparison of pulsar and extragalactic RMs in the anticentre direction indicated a further reversal at larger Galactocentric radius. Rand & Lyne (1994) added more pulsar RMs toward the inner Galactic Plane, and found evidence for two reversals, one about 0.4 kpc inside the Solar Circle, between the Sun and the Sagittarius arm, and another at a Galactocentric radius of 5.5 kpc (with the Sun at 8.5 kpc), near or within the Scutum arm.

At least three classes of model have been proposed for the global structure of magnetic field in the disc of our Galaxy: concentric rings, an asymmetric spiral (ASS) or a bisymmetric spiral (BSS). Rand & Kulkarni (1989) and

Rand & Lyne (1994) argued that pulsar RMs are consistent with a concentric-ring model for the field. Vallée (1991, 1996) has proposed that the field has an ASS form. Although a pure ASS structure has no reversals, modelling suggests that, depending on the initial conditions, reversals can be present (Poezd et al. 1993) and Vallée has argued that the presence of two field reversals inside the Solar Circle is consistent with the ASS model. Early analyses of the RM distribution of extragalactic radio sources (Simard-Normandin & Kronberg 1980; Sofue & Fujimoto 1983) suggested that the Galactic magnetic field has a BSS form, in which the field direction reverses from arm to arm (Heiles 1995, 1996ab). Such structures have been deduced from radio polarization observations of nearby spiral galaxies, such as M81 (e.g. Krause, Beck & Hummel 1989). A statistical study of RMs of a selected sample of pulsars at distances less than ~ 3 kpc (Han & Qiao 1994, 1996) also supported the BSS model for our Galaxy. The pitch angle^{*} of this large-scale field is about -8° .

The magnetic field in the thick disc or halo of our Galaxy may have a different origin to the disc field. Han et al. (1997) showed that the RM distributions of extragalactic radio sources and pulsars in the inner Galaxy (between $l = 270^\circ$ and 90°) are antisymmetric with respect to the Galactic Plane and the meridian through the Galactic Centre. This antisymmetry is not caused by local perturbations, for example, by Loop I, but results from large-scale toroidal magnetic field structure in the thick disc or halo, which has opposite directions above and below the Galactic plane, as in an A0 dynamo mode (see Fig. 19a of Wielebinski & Krause 1993).

Clearly, further work is needed to determine the field structure of our Galaxy, especially that in the disc, and to clarify whether or not the fields in the disc and in the halo are generated or maintained by different mechanisms and are independent of each other (Sokoloff & Shukurov 1990). Observations of pulsar RMs provide a good opportunity to investigate these issues further. There are about 750 known pulsars but, up to now, only 262 pulsars have known RMs (see updated catalogue of Taylor, Manchester & Lyne 1993; TML93). In any quantitative analysis of the disc field, the RMs of high-latitude pulsars should be excluded, further

^{*} There is much confusion in the literature about the definition of pitch angle. For our Galaxy, various authors (e.g. Georgelin & Georgelin 1976; Simard-Normandin & Kronberg 1980; Dame et al. 1986; Grabelsky et al. 1988; Amaral & Lepine 1997) have defined pitch angles to be positive, whereas about an equal number (e.g. Sofue & Fujimoto 1983; Rand & Kulkarni 1989; Han & Qiao 1994; Vallée 1995; Beck et al. 1996; Heiles 1996a) have defined them to be negative.

We propose the following *physical* definition, in principle applicable to all galaxies: the galactic azimuthal angle ϕ is *defined* to be increasing in the direction of galactic rotation. Logarithmic spirals are then defined by

$$R = R_0 \exp(k\phi),$$

where R is the radial distance and R_0 is a scale radius. The pitch angle is $p = \arctan(k)$. This is *negative* for trailing spirals such as our Galaxy, where R increases with decreasing azimuthal angle ϕ . For our Galaxy, the Galactic angular momentum vector points toward the *south* Galactic pole, and ϕ increases in a clockwise direction when viewed from the north Galactic pole.




Figure 1. RM variations of three pulsars. Data of 1980 are taken from Taylor et al. (1993), unpublished data of Hamilton, McCulloch & Manchester (1981; HMM81), of 1987 from Costa et al. (1991), of 1990 from Qiao et al. (1995), of 1991 from van Ommen et al. (1997), and of 1996 from this paper.

limiting the usable pulsar sample. Furthermore, the paucity of measured RMs in the third and fourth Galactic quadrants causes significant uncertainties in the analysis (Han & Qiao 1994).

Between 1996 August 31 and September 5, polarization observations of 66 southern pulsars were obtained with the ATNF Parkes 64-m telescope, mainly at 436 and 660 MHz (Manchester, Han & Qiao 1998). Most of these pulsars were discovered in the recent Parkes southern pulsar survey (Manchester et al. 1996; Lyne et al. 1998). In Sect. 2 of this paper we present RMs for most of these pulsars. We analyse the high-latitude magnetic features of our Galaxy in Sect. 3 and the magnetic field structure in the thin disc in Sect. 4. Current models are discussed in the light of the observational evidence in Sect. 5, and conclusions are presented in Sect. 6.

2 PULSAR ROTATION MEASURES

Pulsar polarization observations were made in bands centred on 436, 660 or 1500 MHz using the Parkes 64-m telescope of the Australia Telescope National Facility. Dual-channel cryogenic systems receiving orthogonal linear polarizations were used at each frequency. Signals were processed in the Caltech correlator (Navarro 1994), which gives 128 lags in each of four polarization channels and folds data synchronously with the pulsar period in up to 1024 bins per period. The data were transformed to the frequency domain, calibrated and dedispersed to form either 8 or 16 frequency sub-bands and corrected for variations in parallactic angle and ionospheric Faraday rotation. Further observational details can be found in Manchester et al. (1998).

In off-line analysis, data from each observation were summed to form two sets of Stokes-parameter profiles for the upper and lower halves of the bandpass. RMs were then

Table 1. Rotation Measures for 63 pulsars

PSR J	l ($^{\circ}$)	b ($^{\circ}$)	Dist (kpc)	DM*	RM (rad m $^{-2}$)	Err	PSR J	l ($^{\circ}$)	b ($^{\circ}$)	Dist (kpc)	DM*	RM (rad m $^{-2}$)	Err
0108–1431	140.9	–76.8	0.10	2.4	–1	3	1544–5308	327.3	1.3	1.29	35.2	–29	7
0133–6957	297.7	–46.7	>2.42	23.0	7	1	1549–4848	330.5	4.3	1.55	57.0	–15	8
0134–2937	230.3	–80.3	>1.78	21.8	13	2	1557–4258	335.3	8.0	7.54	144.5	–37	2
0255–5304	269.9	–55.3	1.15	15.9	–10	12	1559–4438 ^d	334.5	6.4	1.63	58.8	–7	1
0459–0210	201.4	–25.7	1.30	21.0	18	9	1559–5545	327.2	–2.0	5.04	211.0	–150	20
0520–2553	228.4	–30.5	>3.47	33.8	19	15	1603–5657	326.9	–3.3	8.52	264.1	27	5
0536–7543 ^a	287.2	–30.8	1.05	17.6	28	2	1604–4909	332.2	2.4	3.59	140.8	–16	6
0540–7152	282.1	–31.2	2.69	29.4	43	15	1604–7203	316.7	–14.6	2.56	54.4	22	11
0742–2822 ^b	243.8	–2.4	1.90	73.8	156	5	1622–4332	338.3	4.3	8.07	230.7	140	20
0856–6137	278.6	–10.4	6.52	95.0	–70	18	1648–3256	349.6	7.8	5.30	128.3	–60	15
0857–4424	265.5	0.8	6.51	184.4	–75	20	1651–5222	335.0	–5.2	6.39	179.1	–38	5
0924–5302	274.7	–1.9	5.74	155.1	150	20	1700–3312	351.1	5.5	5.62	166.8	–15	3
0934–5249	275.7	–0.7	2.90	99.4	18	6	1722–3712	350.5	–0.5	2.51	99.5	104	3
0942–5657	279.3	–3.0	5.07	160.8	135	4	1741–3927 ^e	350.6	–4.7	4.75	158.5	180	14
0955–5304	278.3	1.2	4.86	156.9	–97	13	1750–3503	355.3	–4.1	5.28	189.4	173	11
1034–3224	272.1	22.1	>4.68	50.9	–8	1	1751–4657 ^f	345.0	–10.2	1.03	20.3	19	1
1036–4926	281.5	7.7	8.39	135.0	–11	6	1801–0357	23.6	9.3	6.66	117.6	32	11
1042–5521	285.2	3.0	6.95	306.0	155	5	1808–0813	20.6	5.8	5.19	151.0	73	7
1116–4122 ^c	284.5	18.1	2.77	40.5	31	16	1829–1751 ^g	14.6	–3.4	5.52	217.8	306	6
1123–4844	288.3	11.6	>8.74	92.9	–7	5	1849–0636	26.8	–2.5	3.66	147.6	–35	21
1126–6942	295.6	–8.1	2.07	55.3	–31	4	1852–2610	9.5	–11.9	2.26	56.8	–21	1
1202–5820	296.5	3.9	4.96	145.8	139	3	1900–2600 ^h	10.3	–13.5	1.70	38.1	–13	2
1210–5559	297.1	6.5	11.83	174.0	58	1	1901–0906	26.0	–6.4	2.42	73.0	44	12
1253–5820	303.2	4.6	2.97	101.0	31	5	1932–3655	2.1	–23.6	>4.41	59.9	6	3
1328–4921	309.1	13.1	>7.78	118.0	170	20	1946–2913	11.1	–24.1	>4.30	44.2	8	7
1328–4357	309.9	18.4	2.23	41.4	–41	3	1949–2524	15.2	–23.4	1.32	22.8	–13	8
1359–6038	311.2	1.1	5.91	295.0	33	5	2038–3816	3.9	–36.7	>2.94	33.9	30	15
1418–3921	320.9	20.5	>5.04	61.5	–15	8	2053–7200 ⁱ	321.9	–35.0	1.01	16.9	9	4
1440–6344	314.6	–3.4	4.03	124.2	29	4	2108–3429	9.7	–42.2	>2.63	30.2	50	20
1527–5552	323.6	0.6	7.07	358.0	34	4	2144–3933	2.8	–49.5	0.18	3.4	–2	10
1527–3931	333.1	14.0	1.88	46.8	4	1	2346–0609	83.8	–64.0	>1.96	22.5	–5	1
1542–5034	328.6	3.6	2.54	95.0	–70	8							

* DM values in cm $^{-3}$ pc. ^aPSR B0538–75: RM= 24 ± 2 by Costa, McCulloch & Hamilton (1991), RM= 21.4 ± 0.5 by Qiao et al. (1995), and RM= 25.1 ± 0.5 by van Ommen et al. (1997); ^bPSR B0742–28: RM= 150.4 ± 0.1 in TML93 from Hamilton, McCulloch & Manchester, unpublished data (HMM81); ^cPSR B1114–41: RM= 50 ± 40 by van Ommen et al. (1997); ^dPSR B1556–44: RM= -2.6 ± 0.3 in TML93 from HMM81, and RM= -5.1 ± 0.5 by van Ommen et al. (1997); ^ePSR B1737–39: RM= 221 ± 29 by van Ommen et al. (1997); ^fPSR B1747–46: RM= 18 ± 3 in TML93 from HMM81 ^gPSR B1826–17: RM= 317 ± 16 by Hamilton & Lyne (1987); ^hPSR B1857–26: RM= -7.3 ± 0.8 in TML93 from HMM81, and RM= -9.5 ± 0.5 by van Ommen et al (1997); ⁱPSR B2048–72: RM= 17 ± 1 by Qiao et al. (1995).

determined by taking weighted means of position-angle differences for bins where the uncertainty was less than 10° ; RM uncertainties were computed from the scatter in position angle differences. For highly dispersed and weakly polarized profiles, we searched the sub-band data in RM for a peak in the linearly polarized intensity $L = (Q^2 + U^2)^{1/2}$. If a significant peak was found, the sub-band data were summed with correction for the Faraday rotation to form the upper and lower bandpass profiles, and a final value of the RM and its error were determined from the position-angle differences and uncertainties as described above.

Rotation measures for 63 pulsars determined in this way are listed in Table 1. The values for Galactic longitude and latitude, distance estimate and dispersion measure in Table 1 are taken from the updated TML93 catalogue. Previously determined RMs for nine pulsars are indicated by

footnotes to the table. For four of these, PSRs J0742–2822, J1116–4122, J1741–3927, J1751–4657 and J1829–1751, the present RMs are either consistent with the old ones, or more precise. The RM of PSR J2053–7200 seems marginally changed from RM= 17 ± 1 rad m $^{-2}$ in 1990 September to RM= 9 ± 4 rad m $^{-2}$ in 1996 September. As shown in Fig. 1 the RMs of PSRs J0536–7542, J1559–4438 (see also van Ommen et al. 1997) and J1900–2600 have varied significantly over several years. Similar RM variations were seen in the Vela (Hamilton et al. 1977; Hamilton, Hall & Costa 1985) and Crab (Rankin et al. 1988) pulsars. These variations are probably caused by a magnetized cloud moving into or out of the line of sight (Hamilton et al. 1985).

Recently, Rand & Lyne (1994) and Qiao et al. (1995) also published a number of new pulsar RMs. Manchester & Johnston (1995) and Navarro et al. (1997) report the RMs




Figure 3. Average field strength along the line of sight versus distance for pulsars with $270^\circ < l < 90^\circ$ and $15^\circ < |b| < 45^\circ$.

of PSR B1259-63 and PSR J0437-4715, respectively. Combining our new measurements with all previously published RMs, we have a total of 318 pulsar RMs.

3 MAGNETIC FIELD FEATURES AT HIGH GALACTIC LATITUDES

3.1 The antisymmetric RM sky

In Fig. 2(a), we show the Galactic distribution of RMs of all pulsars with known RMs and at latitudes $|b| > 8^\circ$. The latitude selection is made to emphasize the magnetic field in the thick disc or halo. Fig. 2(b) shows the filtered RM distribution of extragalactic radio sources (EGRS) of Han et al. (1997), in which RMs discrepant from their nearby neighbours by more than three standard deviations are omitted to more clearly show the large-scale structure of the Galactic field. The antisymmetric distribution is clearly evident in the inner Galactic quadrants. With a few exceptions, the new pulsar RM data shown in Fig. 2(a) are consistent with this antisymmetric RM distribution. Separating the pulsar RM data into three groups at different distances (< 2 , $2 - 4$, and > 4 kpc) shows that the antisymmetry persists, and nearby pulsars show it at higher latitudes. These facts, together with the antisymmetry seen in the RM distribution of EGRS, suggests that the azimuthal fields with opposite directions above and below the Galactic plane are pervasive in the thick disc or halo of our Galaxy. One discrepant region, seen in both pulsar and EGRS RMs, is near $l \sim 310^\circ$, $b \sim 10^\circ$, where positive RMs are found. This region is discussed in Sect. 4.4.

With a few exceptions, both pulsar and EGRS RM data in the region $0^\circ < l < 8^\circ$ have the same sign as those in the fourth Galactic quadrant, $l > 270^\circ$. These sources extend over both positive and negative latitudes and over a range of distances, suggesting a large-scale effect. This result may just reflect field irregularities in this region, but if confirmed to be large-scale, it would suggest that the magnetic field in the halo has a spiral component. If so, the indicated pitch angle is $+8^\circ$, *opposite* in sign to that of fields in the Galactic disc. More RMs of distant pulsars and EGRS around this Galactic longitude are needed to verify this intriguing result.

To estimate the strength of the large-scale magnetic

Table 2. Parameters of pulsars in the south polar region

PSR J	l ($^\circ$)	b ($^\circ$)	Dist (kpc)	RM (rad m $^{-2}$)	$\langle B_{ } \rangle$ (μ G)
0108-1431	140.9	-76.8	0.10	-1 ± 3	-0.7 ± 2.0
2330-2005	49.4	-70.2	0.49	9.5 ± 0.2	1.4 ± 0.3
0034-0721	110.4	-69.8	0.68	9.8 ± 0.2	1.1 ± 0.1
0152-1637	179.3	-72.5	0.79	2 ± 1	0.2 ± 0.1
0206-4028	258.6	-69.6	0.88	-4 ± 5	-0.4 ± 0.5
0134-2937	230.2	-80.2	1.78	13 ± 2	0.7 ± 0.1
0151-0635	160.4	-65.0	1.93	2 ± 4	0.1 ± 0.2
2346-0609	83.9	-64.1	1.96	-5 ± 1	-0.3 ± 0.1

field in the inner quadrants of the Galactic halo, we plot in Fig. 3 the average field strength along the line of sight versus distance for pulsars with $15^\circ < |b| < 45^\circ$. Allowing for the effects of irregular fields and curvature of the field on scales larger than $2 - 3$ kpc, this figure suggests that the large-scale field in the halo has a strength of $1 - 2 \mu$ G.

3.2 A radial field reversal around $l = 225^\circ$, $20^\circ < b < 50^\circ$?

In the region around $l = 225^\circ$ and $20^\circ < b < 50^\circ$, outlined in Fig. 2(a) and Fig. 2(b), pulsar RMs are positive whereas the filtered EGRS RMs are negative. The pulsars in this region are mostly at distances of $1 \sim 2$ kpc, and so the mean field over this path length is directed toward us. However, the mean field through the entire Galactic halo in this direction is directed away from us on average. It appears that there is a reversal of (at least the radial component of) the field at a distance of at least a few kpc. Although more RMs in this region are desirable to confirm this reversal, it demonstrates the large extent of the magnetohalo of our Galaxy and possible field structure within it. Recently, Haffner et al. (1998) have shown that very long H_α filaments exist in this region. If these features are associated, it would imply that the filaments are magnetized.

3.3 Vertical component of local magnetic field

The strength of the magnetic field component perpendicular to the Galactic Plane is a very important parameter since it affects the maximum particle energy in the Galactic halo and hence theories for the origin of cosmic rays (Biermann 1997). It is also an important indicator of field parity in dynamo models (Zweibel & Heiles 1997). Han & Qiao (1994) used the RMs of EGRS at the two Galactic pole regions to investigate this ‘vertical’ component of the local magnetic field. They concluded that the local vertical field B_V has a strength of $0.2 - 0.3 \mu$ G and is directed toward the north Galactic pole. Because of its importance and the difficulty in measuring it in other galaxies, we redetermine B_V based on the filtered RM data for EGRS (Han et al. 1997) and on pulsar RMs in the Galactic pole regions.

We take sources with $|b| > 60^\circ$ and compute the mean vertical component of the field assuming the Taylor & Cordes (1993) model for the Galactic electron distribution. To eliminate any possible contamination by the fields of the North Polar Spur, we do not use RMs of EGRS in the region of $60^\circ < b < 80^\circ$ and $40^\circ < l < 280^\circ$, where the North




Figure 2. (a). Galactic distribution of RMs for pulsars with $|b| > 8^\circ$ from Taylor, Manchester & Lyne (1993), Rand & Lyne (1994), Qiao et al. (1995), Navarro et al. (1997) and this work. The circles represent previously published RMs, and squares indicate RMs from this work. Filled symbols represent positive RMs and the area of the symbols is proportional to $|RM|$ within limits of 5 and 150 rad m^{-2} . The dotted line indicates the antisymmetry with respect to $l = 8^\circ$. The area around $l \sim 225^\circ$ and $b \sim 30^\circ$ outlined by dotted lines is discussed in Sect. 3.3.




Figure 2. (b). Galactic distribution of RMs of selected extragalactic radio sources (EGRS) after Han et al. (1997). Symbols are plotted as in Fig. 2(a).

Polar Spur (Loop I: Berkhuijsen et al. 1971) goes over the polar region (Sofue 1994). A model fit (cf. Sect. 4.3 of Han & Qiao 1994) to $\langle B_{||} \rangle$ values for the remaining 53 EGRS gives a value for the vertical component of $B_V = -0.37 \pm 0.14 \mu\text{G}$. The minus sign means that the vertical field is directed toward the north Galactic pole. The RM contribution of this component is about 3 rad m^{-2} toward the southern pole and -3 rad m^{-2} toward the northern pole. If we subtract this contribution, the RMs of EGRS in the polar regions, mainly

intrinsic or from the intergalactic medium, have a variance of 11 rad m^{-2} .

There are few pulsar RMs in the north polar region, but we can use the RMs of eight pulsars in the south polar region ($b < -60^\circ$) as an independent check of our result. In Table 2, we list the parameters of these pulsars, sorted by distance. We see that four of the five well-determined RMs (error less than $|RM|$) are positive, which is qualitatively consistent with our conclusion from the EGRS.




Figure 4. The RM distribution of pulsars with $|b| < 8^\circ$, projected on the Galactic Plane. The size of the symbols is proportional to the square root of the RM, with limits of 5 and 250 rad m^{-2} with positive RMs represented by a cross and negative RMs by a circle. New measurements are indicated by $\times$ and open squares. The directions of the prevailing Galactic magnetic field, as revealed by pulsar RMs, are indicated by arrows. The approximate locations of four spiral arms are indicated; from the centre out they are, respectively, the Norma arm, the Crux-Scutum arm, the Carina-Sagittarius arm and the Perseus arm (Georgelin & Georgelin 1976; Downes et al. 1980; Caswell & Haynes 1987). The short line at $l = 352^\circ$ separates RMs of opposite sign as discussed in the text, and the dotted circle has a radius of 3 kpc.

4 MAGNETIC FEATURES IN THE GALACTIC DISC

Fig. 4 shows the RM distribution of 189 low-latitude pulsars ($|b| < 8^\circ$), projected onto the Galactic Plane. Separately plotting the data for pulsars above and below the Galactic plane shows little significant difference. In this section, we consider the structure of the local field using RMs of pulsars within 3 kpc of the Sun, and then discuss the evidence for reversals in the field at greater distances.

4.1 The local field

Most previous analyses of pulsar RMs (e.g. Manchester 1974; Lyne & Smith 1989; Rand & Kulkarni 1989) have concluded that the local field is directed toward $l \approx 95^\circ$, i.e., with a pitch angle of $\sim 5^\circ$. This is not consistent with spiral models in which the field lines have a negative pitch angle similar to that of the spiral arms. This discrepancy may be partly explained by selection effects (cf. Han & Qiao 1994), especially the lack of pulsar observations in the fourth (southern) quadrant, and by the influence of the large negative RMs at $l \sim 110^\circ$ to 120° . This group of large negative RMs may be related to the stellar complex in the Perseus arm at these longitudes (e.g. Fig. 2 of Efremov 1995). Passage through a dense ionized region could result in both the large DMs

and the large RMs of these pulsars; the distances of these pulsars may be over-estimated.

All pulsars with distances less than 3 kpc in the region $352^\circ < l < 70^\circ$ (indicated by a short line in Fig. 4) have positive RMs, in contrast to negative RMs of $270^\circ < 350^\circ$, consistent with a pitch angle for the local field of $\sim -8^\circ$. Near and outside the solar circle, the nearby pulsars indicate a regular field running in the clockwise direction whereas those closer to the Galactic Centre indicate a counter-clockwise field.

4.2 Field reversals in the Perseus arm and beyond

Evidence for field reversals in the outer Galaxy (i.e. in or about the Perseus arm) was previously found by Lyne & Smith (1989) and Clegg et al. (1992). Lyne & Smith (1989) noticed that in the region of $105^\circ < l < 135^\circ$ the RM values of extragalactic radio sources are notably smaller than those for most distant pulsars and that for some the signs are reversed, indicating a field reversal beyond the region occupied by the pulsars. Clegg et al. (1992) concluded from the RMs of extragalactic radio sources that there may be a field reversal exterior to the Solar Circle.

As shown in Fig. 4, all distant pulsars, i.e. those outside the Perseus arm, in directions between $l = 55^\circ$ and 80° , have positive RMs. This contrasts with the negative RMs




Figure 5. RMs of pulsars and extragalactic radio sources (a), for $l = 73^\circ \pm 5^\circ$, and (b), for $l = 113^\circ \pm 6^\circ$. Uncertainties in the RMs and distances are indicated for each pulsar. Extragalactic radio sources are plotted at 30 kpc. General trends in the RM variations with distance are indicated by dashed lines.

of nearby pulsars in these directions. Following Clegg et al. (1992) and Rand & Lyne (1994), we plot in Fig. 5(a) RM versus distance for low-latitude pulsars and extragalactic radio sources within 5° of $l = 73^\circ$. The RM increase from negative values of nearby pulsars to the positive ones of these distant pulsars clearly indicates a reversal in the mean field direction in or around the Perseus arm, occurring at a distance ~ 5 kpc. However, the RMs of the EGRS are again negative, indicating a further reversal at distances $\gtrsim 15$ kpc. Clearly, the magnetoionic disc of our Galaxy is very extended (cf. Han & Qiao 1994; Clegg et al. 1992).

RMs in directions around $l \sim 113^\circ$ also give some indication of a field reversal at a distance of about 5 – 7 kpc (but note that the pulsar distances may be overestimated as mentioned in Sect. 4.1). As shown in Fig. 5(b), the RMs of pulsars and EGRS beyond this distance become less negative in general, indicating a positive contribution to the RM from the reversed field in the region beyond the Perseus arm.

Therefore, both of these figures indicate a counter-clockwise field in the region in and beyond the Perseus arm (viewed from the north), in contrast to the clockwise field between the Perseus and Sagittarius arms. Positive RMs around $l \sim 150^\circ$ are consistent with this reversed field. However, in the direction of $180^\circ < l < 190^\circ$, RMs are scattered both positive and negative. In this direction, the lines of sight are approximately perpendicular to the large-scale field, allowing smaller scale irregularities to dominate.




Figure 6. Variation of pulsar RMs with distance in the direction $l = 327^\circ \pm 5^\circ$. Uncertainties are indicated. Trends in the RM variations are indicated by the dashed lines.

4.3 Evidence for a third field reversal inside the solar circle?

Previous work has identified two field reversals inside the Solar Circle, in the Carina-Sagittarius arm and in the Crux-Scutum arm (e.g. Vallée 1991; Rand & Kulkarni 1989; Rand & Lyne 1994). A third field reversal in the Norma arm is predicted by the BSS models of Sofue & Fujimoto (1983) and Han & Qiao (1994). Considering the spiral form of the Norma arm, we choose directions around $l \sim 327^\circ$ and $l \sim 10^\circ$ to examine the RM variations with distance. If the magnetic field follows the spiral arms, as has been seen in nearby spiral galaxies (Beck et al. 1996; Kronberg 1994), it would be almost parallel to the line of sight and hence have the largest RM contribution in these directions.

For $l \sim 327^\circ$ (Fig. 6), the local field within 1 kpc is responsible for positive RMs of the nearby pulsars, indicating a clockwise field near the Sun. Between ~ 1 and ~ 4 kpc, between the Carina-Sagittarius arm and the Crux-Scutum arms, the field direction is reversed. Beyond this distance, the RMs show an increasing trend, indicating a second reversal to a clockwise field between the Crux-Scutum and Norma arms. Beyond this, at distances $\gtrsim 6$ kpc, there is some indication that the RMs are decreasing again, suggesting a possible third reversal in or near the Norma arm.

4.4 Positive RMs in the fourth Galactic quadrant

At longitudes between 280° and 310° and distances greater than 3 kpc, there is a group of pulsars with positive RMs (Fig. 4). These pulsars lie between the Carina and Crux arms and, according to the BSS model, should have negative RMs to complement the positive RMs at longitudes between 35° and 55° . As shown in Fig. 7, these RMs are consistent with the concentric model, but the many positive RMs in the range 300 to 320° at distances greater than 2.5 kpc are not.

Fig. 8 shows that the positive RMs and hence positive $\langle B_{||} \rangle$ at low latitudes in the region between 280° and 310° are mostly above the Galactic Plane, whereas those with negative $\langle B_{||} \rangle$ mostly lie below the Plane. This reversal from the general pattern is probably related to an anomalous region in this direction at distances greater than about 3 kpc. It may represent a large-scale component of the random field or a systematic distortion of the uniform field, and should be investigated in future with more pulsar RM data. Neither




Figure 7. Distribution of pulsar RMs and the concentric-ring model sketched according to Rand & Kulkarni (1989) and Rand & Lyne (1994). Field directions expected for this model are indicated by arrows and the dashed circles show where the field lines reverse. The dotted circles indicate equal distances from the Sun.




Figure 8. Distribution of $\langle B_{||} \rangle$ of all measured pulsar RMs in Galactic coordinates. Plus signs indicate the average field is directed towards us, and circles indicate the average field is directed away from us. The size of symbols is proportional to the field strength within limits of $0.8\mu\text{G}$ and $2.5\mu\text{G}$.

Loop I nor the Gum Nebula can account for it as both of these are much closer.

5 DISCUSSION

The results presented in Sections 3 and 4 show that the magnetic field in our Galaxy has a very complicated struc-

ture. In the thick disc or halo, there is an azimuthal or spiral field with odd symmetry with respect to Galactic plane. The vertical field in the nearby halo is another discriminator for odd parity (Zweibel & Heiles 1997). There is some evidence for such a vertical field in both the pulsar and EGRS data, but at present its statistical significance is weak. In the disc, the field structure is probably related to the spiral arms, as is seen in nearby galaxies (see Beck et al. 1996). Dis-

crimination between the three main models for the disc field proposed so far, the concentric-ring model, the axisymmetric spiral (ASS) model, and the bisymmetric spiral (BSS) model, relies mainly on estimates of the pitch angle of the large-scale field component and the presence and number of field reversals as a function of Galactocentric radius. Unfortunately, the dominance of irregular structures in the field (Ohno & Shibata 1993) makes discrimination between the various models difficult.

Only the concentric-ring model has a pitch angle of zero; the two spiral models predict a pitch angle close to that of the spiral arms. In our Galaxy, both starlight polarization (Heiles 1996a) and the Faraday rotation measurements (Han & Qiao 1993, 1994; Sect. 4 above) suggest a pitch angle for the local field of $\sim -8^\circ$, somewhat less than the pitch angle of spiral arms which is between -8° and -14° , with a best estimate of -12° (Vallée 1995). As discussed in Sect. 4.4, many of the observed RMs, in particular, those around $l \sim 310^\circ$, are not consistent with the ring model. Similarly, the group of positive RMs at large distances around $l \sim 70^\circ$ are inconsistent with this model. Furthermore, there are theoretical difficulties with a circular field structure (Beck et al. 1996). We therefore consider that there is little support for the concentric-ring model of the Galactic magnetic field.

In its pure form, an ASS field has no reversals as a function of Galactocentric radius. Such a field structure can be generated by dynamo action. However, in one-dimensional simulations, Poezd et al. (1993) showed that the ASS field generated by dynamo models could have field reversals. The form and number of these is dependent on assumptions about the initial conditions and the number of galactic rotations and these results need to be confirmed with more realistic (3D) simulations. In external galaxies, field reversals associated with spiral arms have been observed only in BSS candidates (e.g. Beck et al. 1996; Zweibel & Heiles 1997).

In our Galaxy, there is good evidence for one and weaker evidence for two reversals in and beyond the Perseus arm (see also Heiles 1995, 1996ab). Inside the solar Circle, two reversals seem to be more or less certain and we give marginal evidence for a possible third reversal associated with the Norma arm. This means that in all there are probably three, maybe up to five, reversals in the direction of the large-scale Galactic field as a function of Galactocentric radius. This number of reversals is less consistent with ASS models, which have no intrinsic reversals, than with BSS models which have reversals associated with the spiral arms.

On the other hand, the observed antisymmetry of high-latitude RMs suggests that the field in the thick disc and halo of the Galaxy has an A0 form and hence probably generated or maintained by a large-scale dynamo (Han et al. 1997). Yet there is good evidence for close connections between the disc and halo field. For example, around $l \sim 45^\circ$, there are positive RMs extending from $b \sim 0^\circ$ to high positive latitudes (Fig. 8). At high latitudes, these positive RMs form part of the antisymmetric halo field. Many of the disc pulsars with positive RMs are at distances of several kpc (Fig. 4), and their existence is a large part of the argument for the field reversal inside the Sagittarius arm.

Clearly, the question of the origin and structure of the Galactic magnetic field has not yet been answered. There is evidence for ordered magnetic fields in early epochs of

galaxy formation (Udomparsert et al. 1997; Athreya et al. 1998), but dynamo processes (Parker 1997; Kulsrud 1997) will very likely amplify and modify the field structure. More realistic simulations (e.g. Bykov et al. 1997; Rohde & Elstner 1998) show wide variations in the form of the large scale field. Furthermore, the field structure can be complicated by spiral streaming (Lou & Fan 1998; Moss 1998), stellar wind bubbles and supernova events. It is not surprising that the observed field structure in galactic disks cannot yet be fully explained by dynamo models (see the reviews by Kronberg 1994 and Beck et al. 1996).

6 CONCLUSIONS

In this paper, we present RMs for 63 pulsars, 54 of which have no previously published RM data. RMs of three pulsars are shown to have significant variations over timescales of several years. These results reveal the complicated structure of the Galactic magnetic field.

At high Galactic latitudes, the results are largely consistent with the antisymmetric field pattern found by Han et al. (1997). Marginal evidence was found for a leading spiral component in the halo field. An interesting region of reversed radial field at distances of a few kpc was found in the region $l \sim 225^\circ$, $20^\circ < b < 50^\circ$.

Previous observations of fields in the Galactic disc have provided good evidence for two reversals in the large-scale azimuthal field, one inside the Carina-Sagittarius arm at a Galactocentric radius $R \sim 8$ kpc ($R_\odot = 8.5$ kpc), and the other near the Crux-Scutum arm at $R \sim 5.5$ kpc, and weaker evidence for a reversal near or beyond the Perseus arm at $R \gtrsim 10$ kpc (see Heiles 1995, 1996b). The new results strengthen the evidence for the Perseus arm reversal and provide marginal evidence for two additional reversals, one beyond the Perseus arm and one in the inner Galaxy near the Norma arm at $R \sim 3$ kpc. These results, coupled with good evidence that the large-scale disc fields are spiral with a pitch angle $\sim -8^\circ$, are most consistent with a bisymmetric spiral (BSS) form for the disc field. However, positive RMs observed for pulsars located between the Carina-Sagittarius and Crux-Scutum arms in the fourth Galactic quadrant cannot be explained by any spiral model. There is some evidence that the disc and halo fields are connected, further complicating their interpretation.

ACKNOWLEDGMENTS

We thank Drs Elly M. Berkhuijsen, Rainer Beck and the referees for their careful reading of the manuscript and their valuable comments. JLH thanks the Su Shu Huang Astrophysics Research Foundation of CAS and the Director Foundation of BAO for financial support for travel to the ATNF. He also acknowledges the hospitality from Prof. R. Wielebinski during his stay at Max-Planck-Institut für Radioastronomie and financial support from the National Natural Science Foundation (NNSF) of China, the Research Foundation from the Astronomical Committee of CAS. GJQ thanks the NNSF of China for support for his visits at ATNF and acknowledges the financial support from the Climbing

Project – the National Key Project for Fundamental Research of China. JLH and GJQ also thank the ATNF for its hospitality and acknowledge support from the Bilateral Science and Technology Program of the Australia Department of Industry, Science and Tourism. The Parkes telescope is part of the Australia Telescope which is funded by the Commonwealth of Australia for operation as a National Facility by CSIRO.

REFERENCES

- Amaral L. H., Lepine J. R. D., 1997, *MNRAS*, 286, 885
 Athreya R. M., Kapahi V. K., McCarthy P. J., van Breugel W., 1997, *A&A* 329, 809
 Beck R., Brandenburg A., Moss D., Shukurov A. M., Sokoloff D. D., 1996, *ARA&A*, 34, 155
 Berkhuijsen E. M., Haslam C. G. T., Salter C. J., 1971, *A&A*, 14, 252
 Biermann, P., 1997, *J. Phys. G: Nucl. Part. Phys.*, 23, 1
 Bykov A., Popov V., Shukurov A., Sokoloff D., 1997, *MNRAS* 292, 1
 Caswell J.L., Haynes R.F., 1987, *A&A* 171, 261
 Clegg A. W., Cordes J. M., Simonetti J. H., Kulkarni S. R., 1992, *ApJ* 386, 143
 Costa M. E., McCulloch P. M., Hamilton P.A., 1991, *MNRAS* 252, 13
 Dame T. M., Elmegreen, B. G., Cohen, R. S., Thaddeus, P., 1986, *ApJ* 305, 892
 Downes D., Wilson T.L., Bieging J., Wink J., 1980, *A&AS* 40, 379
 Efremov Yu E., 1995, in: *The Formation of the Milky Way*, E. J. Alfaro, A. J. Delgado (eds.), Cambridge, p.46
 Gardner, F. F., Morris, D., Whiteoak, J. B., 1969, *Austral. J. Phys.* 22, 813
 Georgelin Y.M., Georgelin Y.P., 1976, *A&A* 49, 57
 Grabelsky D.A., Cohen R.S., Bronfman L., Thaddeus P., 1988, *ApJ* 331, 181.
 Hamilton P. N., Lyne A. G., 1987, *MNRAS* 224, 1073
 Haffner L.M., Reynold R.J., Tuft S.L., 1998, *ApJ* 501, L83
 Hamilton P. A., Hall P. J., Costa M.E., 1985, *MNRAS* 214, 5p
 Hamilton P. A., McCulloch P.M., Manchester R.N., Ables J.G., Komesaroff M.M., 1977, *Nat.* 265, 224
 Han J.L., Qiao G.J., 1993, in: *Cosmic Dynamo*, IAU Symp.157, Krause F., Rädler K.-H. Rüdiger G. (eds), Kluwer, p.279
 Han J. L., Qiao G. J. 1994, *A&A*, 288, 759
 Han J. L., Qiao G. J. 1996, in: *Pulsars: Problem and Progress*, IAU Collo. 160, Johnston S., Walker M.A. Bailes M. (eds), ASP Conf. Ser. Vol.105, p.485
 Han J. L., Manchester R. N., Berkhuijsen E. M., Beck R., 1997, *A&A* 332, 98.
 Heiles C., 1995, in: *The Physics of the Interstellar Medium and Intergalactic Medium*, Ferrara A. et al., (eds), ASP Conf. Ser. Vol.80, p.507
 Heiles C., 1996a, *ApJ* 462, 316
 Heiles C., 1996b, in: *Polarimetry of the Interstellar Medium*, Roberge W.G. & Whittet D.C.B., (eds), ASP Conf. Ser. Vol.97, p.457
 Krause M., Beck R., Hummel E., 1989, *A&A* 217, 17
 Kronberg P.P., 1994, *Rep. Prog. Phys.* 57, 325
 Kulsrud R.M., 1997, in: *Critical Dialogues in Cosmology*, N. Tuok (ed.), World Scientific, p.328
 Lou Y.-Q., Fan Z., 1998, *ApJ* 493, 102
 Lyne A. G., Smith F. G., 1989, *MNRAS*, 237, 533
 Lyne A. G., Manchester R. N., Lorimer D. R., Bailes M., D'Amico N., Tauris T.M., Johnston S., Bell J.F., Nicastro L., 1998, *MNRAS*, 295, 743
 Manchester R. N., 1971, *ApJS*, 23, 283
 Manchester R. N., 1974, *ApJ*, 188, 637
 Manchester R. N., Han, J. L., Qiao, G. J., 1998, *MNRAS*, 295, 280
 Manchester R. N., Johnston S., 1995, *ApJ*, 441, L65
 Manchester R. N., Lyne A. G., D'Amico N., Bailes M., Johnston S., Lorimer D. R., Harrison P. A., Nicastro L., Bell J. F., 1996, *MNRAS*, 279, 1235
 Mathewson D.S., Ford V.L., 1970, *MmRAS* 74, 139
 Moss D., 1998, *MNRAS* 297, 860
 Navarro J., 1994, PhD thesis, California Institute of Technology
 Navarro J., Manchester R.N., Sandhu J.S., Kulkarni S., Bailes M., 1997, *ApJ*, 486, 1019
 Ohno H., Shibata S., 1993, *MNRAS*, 262, 953
 Parker E.N., 1997, in: *Critical Dialogues in Cosmology*, N. Tuok (ed.), World Scientific, p.309
 Poezd A., Shukurov A., Sokoloff D., 1993, *MNRAS* 264, 285
 Qiao G. J., Manchester R. N., Lyne A. G., Gould D. M., 1995, *MNRAS*, 274, 572
 Rand, R. J., Kulkarni, S. R., 1989, *ApJ*, 343, 760
 Rand R. J., Lyne A. G., 1994, *MNRAS*, 268, 497
 Rankin J.M., Campbell D.B., Isaacman R.B., Payne R.R., 1988, *A&A* 202, 166
 Rohde R., Elstner D., 1998, *A&A* 333, 27
 Sanders W.T., 1995, in: *The Physics of the Interstellar Medium and Intergalactic Medium*, (eds.) Ferrara A., McKee C.F., Heiles C., Shapiro P.R., ASP Conf. Ser. Vol.80, p.498
 Sofue Y., 1994, *ApJ* 431, L91
 Sofue Y., Fujimoto M., 1983, *ApJ* 265, 722
 Sokoloff D. D., Shukurov A., 1990, *Nat.*, 347, 51
 Simard-Normandin M., Kronberg P. P., 1980, *ApJ*, 242, 74
 Taylor J. H., Cordes J. M., 1993, *ApJS* 411, 674
 Taylor J. H., Manchester R. N., Lyne A. G., 1993, *ApJS*, 88, 529 (TML93)
 Thomson R. C., Nelson A. H., 1980, *MNRAS*, 191, 863
 Udomprasert P. S., Taylor G. B., Pearson T. J., Roberts D. H., 1997, *ApJ* 483, L9
 Vallée J. P., 1991, *ApJ*, 366, 450
 Vallée J. P., 1993, *ApJ*, 419, 670
 Vallée J. P., 1995, *ApJ*, 454, 119
 Vallée J. P., 1996, *A&A*, 308, 433
 van Ommen, T. D., D'Alessandro, F. D., Hamilton, P. A., McCulloch, P. M. 1997, *MNRAS*, 287, 307
 Wielebinski R., Krause F., 1993, *A&AR*, 4, 449
 Zweibel E.G., Heiles C., 1997, *Nat.*, 385, 131